

1 Past and projected climate change impacts on heat-related child
2 mortality in Africa

3 Sarah Chapman^{*1}, Cathryn E Birch¹, John H Marsham¹, Chérie Part², Shakoor Hajat²,
4 Matthew F Chersich³, Kristie L Ebi⁴, Stanley Luchters^{5,6,7}, Britt Nakstad^{8,9}, Sari Kovats²

5 ¹ School of Earth and Environment, University of Leeds, Leeds, United Kingdom

6 ² Centre on Climate Change & Planetary Health, London School of Hygiene & Tropical
7 Medicine, London, United Kingdom

8 ³ Wits Reproductive Health and HIV Institute, Faculty of Health Sciences, University of the
9 Witwatersrand, Johannesburg, South Africa

10 ⁴ Department of Global Health, University of Washington, Seattle, Washington

11 ⁵ Institute of Human Development, Aga Khan University, Nairobi, Kenya

12 ⁶ Department of Public Health and Primary Care, Ghent University, Belgium

13 ⁷ Department of Epidemiology and Preventive Medicine, Monash University, Australia

14 ⁸ Division Paediatric Adolescent Medicine, Institute of Clinical Medicine, University of Oslo,
15 Oslo, Norway

16 ⁹ Department of Paediatrics and Adolescent Health, University of Botswana, Gaborone,
17 Botswana

18 * corresponding author email: earsch@leeds.ac.uk

19

1 **Abstract**

2 Children (<5 years) are highly vulnerable during hot weather due to their reduced ability to
3 thermoregulate. There has been limited quantification of the burden of climate change on
4 health in sub-Saharan Africa, in part due to a lack of evidence on the impacts of weather
5 extremes on mortality and morbidity. Using a linear threshold model of the relationship
6 between daily temperature and child mortality, we estimated the impact of climate change on
7 annual heat-related child deaths for the current (1995–2020) and future time periods (2020–
8 2050). By 2009, heat-related child mortality was double what it would have been without
9 climate change; this outweighed reductions in heat mortality from improvements associated
10 with development. We estimated future burdens of child mortality for three emission
11 scenarios (SSP119, SSP245 and SSP585), and a single scenario of population growth. Under
12 the high emission scenario (SSP585), including changes to population and mortality rates,
13 heat-related child mortality is projected to double by 2049 compared to 2005-2014. If 2050
14 temperature increases were kept within the Paris target of 1.5°C (SSP119 scenario),
15 approximately 4,000 – 6,000 child deaths per year could be avoided in Africa. The estimates
16 of future heat-related mortality include the assumption of the significant population growth
17 projected for Africa, and declines in child mortality consistent with Global Burden of Disease
18 estimates of health improvement. Our findings support the need for urgent mitigation and
19 adaptation measures that are focussed on the health of children.

20 **Keywords:** climate change, health impacts, heat stress, child mortality, Africa

21

22

23 **Introduction**

1 Climate change negatively affects human health through exposure to extreme weather
2 and climate events, including the direct effect of high temperatures on heat stress, and
3 indirect effect of temperature on infectious disease and food safety and security (Sheffield
4 and Landrigan 2011, Rylander et al. 2013). High temperatures are responsible for a
5 significant burden of disease. The global estimate of heat-related mortality across all ages is
6 approximately 500,000 deaths per year (Zhao et al. 2021). Temperatures are already
7 increasing in Africa, by between 0.2 – 0.4°C per decade since 1980 (Trisos et al. 2022). As
8 temperatures continue to rise with climate change, heat deaths are expected to increase.

9 Children under five years old are especially vulnerable to the impacts of heat
10 exposure (Helldén et al. 2021). Very young children have a limited ability to thermoregulate
11 (Sheffield and Landrigan 2011, Rylander et al. 2013, Son et al. 2017). Children in low-
12 income settings with high temperatures are particularly vulnerable to heat due to pre-existing
13 burdens of infection and undernutrition, poor healthcare systems, and dwellings that do not
14 provide sufficient protection from the heat (Sheffield and Landrigan 2011, Rylander et al.
15 2013). In high-income countries, heat-related deaths are largely restricted to deaths in older
16 persons. In low-income settings, high temperatures are shown to increase acute mortality in
17 children and young adults, representing a considerable burden in terms of years of life lost
18 (Sewe et al. 2018). Due to limitations associated with both health and environmental data,
19 there has been little observational research in low-income and middle income countries on
20 heat impacts, particularly in children (Basu et al. 2015, Green et al. 2019, Berrang-Ford et al.
21 2021).

22 Better understanding how child mortality could be affected by climate change is vital
23 for ensuring prioritization of health in national adaptation programmes. The future impacts of
24 climate change on child mortality have previously been quantified for malnutrition and
25 diarrhoeal disease (World Health Organization 2014) but not for heat-related mortality.

1 Countries in Africa are set to undergo major development and demographic changes in the
2 coming decades, with the potential to ameliorate or amplify future climate change impacts.

3 The aim of this study is to estimate the impact of climate change on annual heat-
4 related child deaths for the current (1995–2020) and future time periods (2020–2050) in
5 Africa, using published estimates of the relationship between ambient temperature and acute
6 mortality in selected populations in Africa. First, we undertook an attribution study to
7 quantify heat-related child mortality due to climate change in Africa over the period 1995–
8 2020. We use observed population growth and the reductions in child mortality rate (due to
9 improvements in health services, disease control, and the social determinants of health). We
10 then conduct a comparative risk assessment to project the numbers of heat-related child
11 deaths that occur in future decades (2020–2050) under low, medium, and high emissions
12 scenarios (SSP119, 245, and 585; Meinshausen et al. 2020), considering future population
13 growth and the continued decline in child mortality rates. As there is very limited
14 epidemiologic information available on the relationship between temperature and mortality in
15 children in African countries, we use published coefficients from two studies to represent
16 high and low bounds of heat-related mortality burdens for the continent. This work, therefore,
17 provides indicative estimates of impacts. Despite the obvious limitations in basing estimates
18 for Africa on studies in two locations, we believe providing an indicative first estimate of the
19 scope of the issue is an important step in drawing attention to the problem of heat-related
20 mortality in children, and the need for further research in this area.

21

22 **Methods**

23 We attributed heat deaths in children in Africa to anthropogenic emissions using the
24 DAMIP experiment of CMIP6. This method has been used previously to estimate that a third

1 of all heat-deaths in the period 1991–2018 can be attributed to climate change globally
2 (Vicedo-Cabrera et al. 2021). Attributing observed changes in health outcomes to climate
3 change can be difficult as key health determinants also change over decadal scales (Ebi et al.
4 2017). In this study, we have made a first estimate of the impact of climate change on heat-
5 related child mortality in the present day and future, using CMIP6 climate scenarios,
6 population and mortality projections, and published heat-mortality relationships in Africa.

7 *Description of observed climate change and counterfactual datasets*

8 We attributed heat-mortality to climate change by comparing two climate change
9 scenarios, one corresponding to the historical (factual) climate, and an alternative
10 (counterfactual) that corresponds to a world without anthropogenic climate change. The
11 scenarios are from the DAMIP (Detection and Attribution Climate Model Intercomparison)
12 experiment of CMIP6, designed to distinguish the climate impacts of different forcings
13 (O’Neill et al. 2016). For the factual scenario, we used mean daily air temperature from
14 ‘historical’ climate simulations from 1995 up to 2014, merged with SSP245 for 2015 – 2020,
15 because 2015 is the start of the future simulation. These simulations are driven by natural and
16 anthropogenic forcings; and are designed to represent actual historical climate. The
17 corresponding counterfactual scenario is from the ‘hist-nat’ experiment, where only natural
18 forcings (no anthropogenic greenhouse gases) were considered. Thus, the difference between
19 the ‘hist-nat’ and ‘historical’ experiments quantifies the impact of anthropogenic forcings on
20 the climate in Africa. Thirteen climate models were used in the analysis (see Supplementary
21 Material Table 1).

22 *Description of the present-day population and mortality datasets*

23 We used the gridded (1km² resolution) WorldPop under-5 population data (WorldPop
24 and Center for International Earth Science Information Network (CIESIN) 2018) and national

1 under-5 all-cause mortality rates from UNICEF (UNICEF Data 2019). 2000 is the earliest
 2 year for which age-structured, gridded population data are available, which is why we limited
 3 our historical analysis to 1995 – 2020. The mortality data were available as country totals and
 4 we mapped child mortality values to individual grid cells using the population distribution for
 5 each time point (Figure S1). The mortality data were available as yearly totals and daily all-
 6 cause mortality from equation 1 was calculated as annual all-cause mortality divided by 365.

7 The time periods of the datasets used in the attribution study are shown in Table 1.
 8 For the first time period, population and mortality data from the middle of the period were
 9 used. For the 2015 – 2020 period, we used the most recent population and mortality data
 10 available, which was for 2019.

11 Table 1. Climate, population and mortality data for attribution study.

Data	Time period		
	1995 – 2020 (25 years)		
Climate	1995 – 2004	2005 -2014	2015 - 2020
Population	2000	2010	2019
Mortality	2000	2010	2019

12
 13 *Description of the scenario data*

14 To estimate future heat-related child mortality from 2020 – 2050, we used three
 15 scenarios of climate change (SSP119, SSP245 and SSP585), and one scenario of population
 16 growth and one scenario of child mortality which assumes that the current declines in child
 17 mortality rates will continue. SSP119 is the scenario most likely to achieve the 1.5°C target
 18 under the Paris Agreement; SSP585 is a high fossil-fuel emission scenario; and SSP245 is a
 19 middle-of-the-road emission scenario (Meinshausen et al. 2020). From the CMIP6 climate
 20 models, we obtained daily average temperature for each day between 2020 – 2050. For
 21 population and mortality data, we used information from the middle of the decade (i.e., the
 22 2020 – 2029 period used 2025 population and mortality data).

1 For future populations (see Figure S2), we used the SSP2 (middle-of-the-road)
2 population projection (KC and Lutz 2017). We obtained age-structured country totals from
3 the Wittgenstein Centre (Wittgenstein Centre for Demography and Global Human Capital
4 2018).

5 For future all-cause mortality, we used the Global Burden of Disease (GBD) reference
6 scenario (Foreman et al. 2018, Global Burden of Disease Collaborative Network 2020). We
7 chose this projection because it is an evidence-based and plausible projection of health
8 futures in Africa and because of the large number of modeled causes of death (Foreman et al.
9 2018). The GBD reference scenario tends to give lower future all-cause mortality than the
10 World Health Organization projections. The impact of lower all-cause mortality would be a
11 lower absolute number of heat-related deaths. As the GBD projections only extend to 2040,
12 we extrapolated to 2050 using the trend from 2020 – 2040 and a second-order polynomial
13 (Figure S2). This was applied on a country-by-country basis.

14 Future population and mortality data were distributed within countries according to
15 the distribution of population in 2019. Migration within and between countries was not
16 included due to lack of any demographic scenarios. Africa is likely to undergo significant
17 urbanisation which may in fact increase local temperature exposures for urban populations.
18 Alternatively, large scale movement to cooler climates would lead to an over-estimate of the
19 modelled heat mortality. Overall, future population movement and its effect on heat
20 exposures is highly uncertain.

21 Future population and mortality projections were adjusted using the 2000 and 2010
22 population and mortality data, respectively, and the linear-scaling method (see
23 Supplementary Material for further details).

24 *Description of the health impact model*

1 We quantified the heat-related child (under-5) mortality in each grid cell in each
2 individual climate model using the following equation (Hajat et al. 2014):

$$DAT = \frac{Pop_{fut}}{Pop_{base}} \times \frac{DM}{e^{(b \times T_d - Thres)}} \times (e^{(b \times T_d - Thres)} - 1) \quad (1)$$

3

4 Where DAT = daily attributable deaths due to temperature, DM = daily average mortality
5 from all causes, T_d = daily average temperature, b = coefficient of the exposure-risk
6 relationship, $thres$ = threshold above which temperature mortality impacts occur. Under 5
7 population in the year 2000 was used as the base population (Pop_{base}). Pop_{fut} refers to future
8 under 5 population.

9 The coefficient, b , and the threshold are from two published time-series regression
10 studies in Ghana and Kenya (Azongo et al. 2012, Egondi et al. 2012). The study in Kenya
11 was based on 5 years of data, while the study in Ghana was based on 15 years of data
12 (Azongo et al. 2012, Egondi et al. 2012). Limitations common to all epidemiological studies
13 include missing data, and limited data on time varying confounding factors (Azongo et al.
14 2012, Egondi et al. 2012).

15 Both studies used the 75th percentile of local daily average temperature as the heat
16 threshold above which risk of death increases linearly as temperatures rise; therefore, we
17 assumed the 75th percentile of the temperature distribution unique to each grid cell to model
18 localised heat thresholds. There is some variation in the threshold temperatures at which
19 mortality impacts occur between populations (Hajat and Kosatky 2010). However, applying
20 the same threshold percentile across the region is a reasonable approximation that has been
21 used in other studies (Honda et al. 2014). The 75th percentile is towards the lower end of the
22 percentile thresholds usually used for heat-mortality relationships. A higher threshold would
23 result in a lower absolute number of deaths.

1 Exposure effects were based on lags of 0 and 1 days, which were demonstrated to be
2 the critical periods of exposure for heat in the studied populations (Sheffield et al. 2018,
3 Schinasi et al. 2020).

4 The temperature threshold was calculated from the 1995 – 2010 period of the
5 historical run from each individual model. This period was chosen as it encompasses the time
6 periods used in the epidemiology studies. The work in Ghana and Kenya found that mortality
7 increased linearly by 0.61 and 1.0% respectively for each 1°C increase in daily average
8 temperature above the threshold. In other words, effect estimates were based on linear-
9 threshold models. These values were converted into the coefficient, b , using the following
10 equation:

$$b = \ln\left(\frac{\% \text{ increase per } 1^{\circ}\text{C}}{100} + 1\right) \quad (2)$$

11 Where b represents just the linear portion of the temperature-mortality relationship
12 characterised in the two studies. We calculated heat-related child mortality using the
13 coefficients from both studies, and present the results as a range. There are limitations in
14 using functions from only two locations in Africa, but we have developed this model using
15 the best information available at this time. There are other studies available that examine heat
16 related child mortality in Africa, but the two used here are the only ones that use consistent
17 methods to quantify the acute temperature-mortality relationships in children in Africa. Due
18 to differences in methods, it would not be straightforward to combine information from the
19 other African studies, ie., Scovronick et al. (2018) uses maximum temperatures than than
20 mean and Mrema et al. (2012) uses monthly temperatures rather than daily.

21 Prior to calculating mortality, all climate models were bias corrected (see below), and
22 all input data sets (including population and mortality datasets) were regridded to a 0.5° x 0.5°
23 resolution, the most common resolution of the input CMIP6 models.

1 *Decomposition of mortality into components*

2 We used the method of Das Gupta to decompose the contribution of changes in
3 climate, population, and all-cause mortality rate to changes in heat-related child mortality
4 (1993). To calculate the population, mortality and exposure (climate) effect, we used the
5 following equations (Das Gupta 1993):

$$\text{Population effect: } (p - P) \left[\frac{me + ME}{3} + \frac{mE + Me}{6} \right] \quad (3)$$

$$\text{Mortality effect: } (m - M) \left[\frac{pe + PE}{3} + \frac{pE + Pe}{6} \right] \quad (4)$$

$$\text{Exposure effect: } (e - E) \left[\frac{pm + PM}{3} + \frac{pM + Pm}{6} \right] \quad (5)$$

6 Where p , m and e refer to population ($\frac{Pop_{fut}}{Pop_{base}}$ from equation 1), total all-cause under-5

7 mortality (DM from equation 1), and exposure to high temperatures respectively

8 ($\frac{(e^{(b \times T_d - Thres)} - 1)}{e^{(b \times T_d - Thres)}}$ from equation 1), and the lowercase refers to the historical (baseline) value

9 and upper case refers to the future value.

10 *Bias-correction of climate model data*


11 Daily average temperature from all CMIP6 climate models (historical, hist-nat and
12 future) were bias-corrected using the linear-scaling method with monthly correction factors
13 (Teutschbein and Seibert 2012) and the CRU TS4.0 gridded climate dataset (Harris 2019);
14 1970 – 1989 was the reference period for bias-correction.

15 **Results**

16 *Present day climate change impacts on temperature*

17 The impact of climate change on average temperatures in Africa, 1995 – 2020, is
18 shown in Figure 1. The ensemble mean of the CMIP6 models is cooler for the hist-nat

1 (without anthropogenic climate change) scenario than the historical (with climate change)
 2 scenario. The average temperature increases over time in the historical scenario, while it stays
 3 relatively stable in the hist-nat scenario. The CRU observational data also shows this trend of
 4 increasing temperature over time, and is of a similar magnitude to the increase in temperature
 5 in the CMIP6 ensemble mean. The increasing temperature trends over time in Africa are well
 6 documented in the literature (Trisos et al. 2022).




7

1 Figure 1. Average monthly temperature in Africa from 1995 – 2020, for the CRU
2 observational dataset (row 1), the CMIP6 ensemble mean for the historical scenario (row 2)
3 and the hist-nat scenario (row 3)

4 *Current heat-related child mortality due to climate change*

5 Heat-related child mortality for the historical and hist-nat scenarios for the periods
6 1995 – 2004 and 2011 – 2020 are shown in Figure 2. Between 1995 and 2004, the CMIP6
7 ensemble mean heat-related child mortality in Africa was approximately 7 000 – 11 000
8 deaths per year, depending on whether a lower (coefficient = 0.61) or higher (coefficient =
9 1.0) sensitivity to heat was used. Without climate change, this would have been
10 approximately 4 000 – 6 000 deaths per year. In the 2011 – 2020 decade, with climate
11 change, these numbers increase dramatically to approximately 12 000 – 19 000 deaths per
12 year, depending on whether low or high sensitivity to heat was used. Without climate change,
13 this would have been approximately 5 000 – 8 000 deaths per year. From 2009 onwards, heat-
14 related child mortality with climate change was double what it would have been without
15 climate change. Data for each country in Africa are presented in Supplementary Material
16 Table S2.




1

2 Figure 2. Heat-related child (under-5) mortality from 1995 – 2004 and 2011 – 2020 A)
 3 Annual average mortality in Africa over the decade, B) Annual average mortality per 100 000
 4 children in Africa over the decade. Bars show 5th and 95th percentile from CMIP6 ensemble.
 5 Based on lower sensitivity to heat (coefficient = 0.61).

6 We decomposed the overall change in total heat-related child mortality from 1995 –
 7 2004 to the recent decade of 2011 – 2020 into contributions from population growth,
 8 declining all-cause mortality rates, and climate change (Figure 3A, see Methods for details).
 9 Population growth (Figure 3A) meant that the total number of heat-related child deaths
 10 increased over time, even without climate change (Figure 2, panel A, white bars), and it was
 11 the largest contributor to the increase in total heat-related child mortality. Declining all-cause
 12 mortality rates (Figure 3) mean that without climate change, the rate of heat-related child
 13 mortality (Figure 2, panel B, white bars) would have declined over time. However, these
 14 potential gains were outweighed by the impact of rising temperatures from climate change

1 (Figure 3). These conclusions were the same when assuming a higher sensitivity to heat, with
 2 higher absolute numbers of child deaths (coefficient = 1.0, see Figure S3).

3 Figure 3, Panel B, shows how population growth, declining all-cause mortality rates,
 4 and climate change are projected to contribute to changes in heat-related child mortality in
 5 the 2040s relative to 1995 – 2004. Population growth begins to slow after the 2020s, and in
 6 some countries begins to decline by the 2040s (Figure S4) relative to 2019. Therefore, the net
 7 change in heat-related child mortality is high in the historical scenario (Figure 3, panel A)
 8 compared to the climate change scenarios (Figure 3, panel B).



9

10 Figure 3. Contribution of changes in population, declining all-cause mortality rates, and
 11 climate change to changes in annual average heat-related child mortality. Net change in
 12 annual heat-related child deaths is in grey. 1995–2004 was the base period for calculating


1 change. Panel A) Historical (with climate change) and hist-nat (without climate change)
2 scenarios. Panel B) Climate change scenarios, SSP119, 245 and 585 (compared to historical
3 scenario). Based on lower sensitivity to heat (coefficient = 0.61).

4 *Projected heat-related child mortality with climate change*

5 Figure 4 shows that heat-related child mortality in Africa is projected to increase over
6 the next thirty years as temperatures rise for all scenarios except SSP119. In the 2040s, for
7 the SSP119 scenario, heat-related child mortality begins to stabilize as temperatures stabilize
8 and health care and socio-economic factors improve, as represented by declining all-cause
9 mortality rates (Figures 3 and 4). Under SSP245 and SSP585, total heat-related child
10 mortality continues to rise as temperatures continue to rise. When comparing these scenarios
11 with the latest decade from the historical scenario (2005 – 2014), by the 2030s, annual
12 average total heat-related child mortality in Africa increases by approximately 48% under
13 SSP119, 53% under SSP245, and 55% under SSP585. By the 2040s, this increase is projected
14 to be 31% for SSP119, 49% for SSP245, and 75% for SSP585. By 2049, under SSP585,
15 annual heat-related child mortality in Africa is projected to double the annual average from
16 2005 – 2014. See Figure S5 for results as a rate rather than total; this shows that even when
17 removing population increase by considering mortality as a rate, large increases in deaths are
18 projected with climate change from the 2040s onwards.

19 Figure 4 shows the benefits of heat deaths avoided by reducing carbon emissions. By
20 the 2030s, following the SSP119 scenario would mean on average approximately 600 – 1000
21 children could be saved per year in Africa compared to SSP585, depending on sensitivity to
22 heat, while SSP245 would mean on average approx. 200 – 300 lives saved compared to
23 SSP585. In the 2040s, on average approx. 4,000 – 6,000 lives could be saved annually by
24 following SSP119 compared to SSP585, and approx. 2,000 – 3,000 by following SSP245

1 compared to SSP585. There is a large spread in the projected temperature increases under
 2 SSP585 by mid-century (Figures 4 and 6). Based on the 5th-95th percentile range from the
 3 CMIP6 ensemble, child mortality could reach over 23,000 deaths per year with low
 4 sensitivity to heat and 38,000 with high sensitivity to heat in 2049 under SSP585. If reality
 5 turned out to be closer to the upper estimates, the benefits to following SSP119 would be
 6 greater.




7
 8 Figure 4. Projected annual heat-related child mortality in Africa with climate change.
 9 Population growth (SSP2 population scenario) and declining all-cause mortality rates (GBD
 10 reference scenario) were assumed., assuming lower sensitivity to heat (coefficient = 0.61).
 11 Panel A) Box and whisker plot for 2015. Panel B) Ensemble mean child mortality from 2015
 12 – 2049. Panel C) Box and whisker plot for 2049. For panels A and C), the box shows 25th –
 13 75th percentile. Box lines shows median. Whiskers show 5th – 95th percentile from CMIP6
 14 ensemble, circles show values beyond this range. Only CMIP6 models with data for all three

1 scenarios were included. When all CMIP6 models were included, 5-95th percentile range is
2 larger.

3 *Heat-related child mortality beyond 2050*

4 We did not project heat-related child mortality beyond 2050 because of uncertainties
5 in the socio-economic projections. Impacts in the next few decades are more relevant for
6 public health and planning horizons. In addition, there is relatively little divergence in
7 projected temperature from the different emissions scenarios until after 2040, and so
8 temperature increases to mid-century need to be planned for regardless.



9

10 Figure 5. Number of days per year above the 1995-2010 75th percentile threshold of daily
11 average temperature, calculated from CMIP6 climate models. Shading shows model range
12 (5th – 95th percentile from CMIP6 ensemble). Vertical line at 2050. Only CMIP6 models with
13 data for all three climate scenarios included.

1 As the century progresses, under SSP585 and SSP245, most days in the year would be
2 above the temperature mortality threshold (Figure 5). African populations would be
3 experiencing an entirely new climate. Already by 2050, many tropical locations are expected
4 to experience climates not currently experienced anywhere on earth (Bastin et al. 2019).
5 Under these circumstances present-day temperature-mortality relationships are likely to be a
6 significant underestimate of the true effects (Rocklöv and Ebi 2012). Uncertainties in the
7 population and mortality projections (discussed below in the Discussion section), also make
8 projecting beyond 2050 challenging.

9 **Discussion**

10 Our results show that climate change is already having a substantial impact on child
11 mortality in Africa. Target 4.a of the Millennium Development Goals was to reduce mortality
12 in under-5s by two-thirds between 1990 and 2015 (World Health Organization 2005). While
13 significant reductions in under-5 mortality have been achieved since 1990, the target of the
14 Millennium Development Goals was not met (United Nations 2015). Heat-related mortality
15 from climate change may have undermined gains made in improved child health. The
16 Sustainable Development Goals have the goal of reducing under-5 mortality by 2030 (United
17 Nations 2020). Our results suggest that if climate change is not kept to 1.5°C of warming,
18 rising temperatures would make meeting the SDG target increasingly difficult.

19 The limitations of this study include the reliance on heat-mortality relationships from
20 existing literature that allowed the use of only two regions. Population-specific temperature
21 mortality relationship have been found to vary by latitude, altitude, socio-economic factors
22 such as income inequality, and factors relating to the build environment, such as prevalence
23 of air-conditioning use (Liu et al. 2021). It is likely that heat-mortality relationships in
24 children to vary across populations in Africa. However, there are only a very small number of

1 other studies on heat-mortality relationships for children in Africa. Due to different methods
2 used, we were unable to combine information from these studies. The benefit of the effect
3 estimates we used for our heat-mortality relationships is the use of a consistent, widely
4 accepted method that makes comparisons more robust and the presentation of results as a
5 range more meaningful. We acknowledge that there is undoubtedly variability in heat
6 vulnerability across Africa due to additional factors we could not model due to a lack of
7 epidemiologic evidence. Furthermore, the heat-mortality relationship may change over time.
8 Therefore we have more confidence in the results from the historical period than the future
9 period. The studies we used from Ghana and Kenya found a 0.61 and 1% increase in
10 mortality for each 1°C rise in temperature above the threshold temperature, respectively. A
11 review of studies found the heat mortality impact is usually between 1 and 3% (Hajat and
12 Kosatky 2010); however, these results are not for children, and only one of these studies was
13 for Africa. If studies on children in Africa report a stronger heat-mortality relationship, then
14 the heat-related child mortality would be higher than estimated here. This, however, would
15 not change the qualitative impact of climate change on mortality, i.e., that there will be
16 greater heat-related child mortality with SSP585 than SSP119, and mitigation will save lives.

17 More research and data, based on comparable methods, are needed to quantify
18 mortality and morbidity outcomes related to heat in low-income settings, and to allow for
19 robust estimation of exposure effects in high-risk groups such as children. Because the data
20 required to estimate temperature-mortality relationships may not be routinely collected in
21 many low-income countries, research should focus on other ways of understanding heat
22 impacts in the absence of local data.

23 We did not assess cold-related mortality due to the limited available information in
24 this region. The cold effect was not statistically significant in the Kenya study and in Ghana
25 was over 12 times smaller than the heat effect. Cold effects occur at longer lag structures and

1 there are some doubts regarding a causal association in some settings (Arbuthnott et al.
2 2018). We found no robust estimates for heat effects on very young children, such as
3 newborns (< 1 month), or infants (1 year), despite these groups being especially vulnerable to
4 heat and cold (Xu et al. 2012).

5 We modelled adaptation as the general improvements in health and not as specific
6 changes to the temperature-mortality function (World Health Organization 2014).
7 Adaptation is often omitted from heat-related mortality assessments of climate change due to
8 the lack of robust estimates about the extent to which adaptation will reduce impacts (Wang
9 et al. 2019). Successful adaptation would lead to a reduced burden of future heat-related
10 mortality. Adaptation options may be limited in this region due to limited economic capacity,
11 specifically the affordability of space cooling. Emerging research in African settings has
12 identified some low costs interventions, including cool roofs, and behaviour change
13 interventions for reducing occupational heat risks (Spencer et al. 2022). This study supports
14 the need for increased prevention of heat-related risks in Africa, including the adoption of
15 public health measures such as heat health action plans. More research is needed on effective
16 measures to reduce heat risk, including behavioural responses that can easily be adopted in
17 low-resource settings (Rocklöv and Ebi 2012).

18 A further cause of uncertainty is with the all-cause mortality rate projections
19 (Foreman et al. 2018) that decline approximately 50% by 2040 (Figure S2), based on
20 assumptions regarding improvements in life expectancy (United Nations 2019), socio-
21 economic factors, and health-care (Foreman et al. 2018). Projections that assume a decline in
22 all-cause mortality by the end of the century may not be consistent with a future world under
23 a high-emission climate change scenario. High temperatures affect the incidence of diseases,
24 such as malaria (Yé et al. 2007, Chirombo et al. 2020), diarrhoea (Smith et al. 2014), and
25 gastrointestinal and respiratory infections (Phung et al. 2015, Kim et al. 2016). These are

1 important causes of mortality and morbidity in children under age 5 (World Health
2 Organization 2005, 2018). Premature birth, which is in the top three causes of child death in
3 Africa and the leading cause worldwide (World Health Organization 2018), has been shown
4 to increase with exposure to high temperatures during pregnancy (Chersich et al. 2020). In
5 addition, climate change is expected to increase extreme events, and in many areas will make
6 accessing clean drinking water more difficult (Smith et al. 2014). It is conceivable therefore,
7 that if temperatures continue to rise, deaths from these causes will also rise (Sheffield and
8 Landrigan 2011). If this is the case, calculations of heat-related child mortality based on
9 assumptions of declining total mortality would be underestimate future heat mortality.

10 Substantial benefits to health can be achieved by reducing carbon emissions. Even
11 without projecting child mortality out to 2100, it is clear from Figure 5 and the preceding
12 analyses that by 2100, substantial health benefits would be achieved by following the SSP119
13 emissions pathway. The current global emissions trajectory tracks the high-end of the
14 emissions scenarios (Friedlingstein et al. 2014, Schwalm et al. 2020). Rapid and effective
15 mitigation and emissions reductions would entail significant reductions in heat-related
16 mortality by the 2040s. The contrast between child mortality in SSP585 and SSP119 by the
17 end of the century would be larger.

18 **Conclusion**

19 Our results show that climate change, through increasing exposure to high
20 temperatures, may have already led to double the heat related child mortality compared to
21 what would have been expected without climate change. This underscores the need for more
22 ambitious mitigation measures to protect vulnerable populations and the need for proactive
23 and effective adaptation. In the absence of ambitious emissions reductions targets and the
24 introduction of adaptation measures, heat stress impacts on child mortality will increase, and

1 under a high emissions scenario may double in Africa by 2049. Studies in other regions with
2 tropical and sub-tropical climates have shown an impact of high temperatures on child
3 mortality (Xu et al. 2012, Son et al. 2017), therefore climate change is likely to have
4 impacted heat-related child mortality burdens in other parts of the world, particularly south
5 Asia and South America (Romanello et al. 2021). Our understanding of heat impacts on
6 children, and how heat-mortality relationships could change with large amounts of warming,
7 is limited due to the lack of physiological and epidemiological studies on heat stress in
8 children, particularly in low-resource settings. More research is urgently needed to
9 understand how some of the most vulnerable members of our population may be impacted by
10 extreme heat and which health interventions would effectively reduce heat impacts. Our
11 results highlight the urgent need for health policy to focus on heat-related child mortality, as
12 our results show it is a serious present-day issue, which will only become more pressing as
13 the climate warms.

14 **Data availability statement**

15 All data analysed in this paper is available publicly. CMIP6 model data available from ESGF
16 (<https://esgf-node.llnl.gov/projects/cmip6/>). Age-structured gridded population data available
17 from WorldPop (<https://www.worldpop.org/geodata/listing?id=65>). Under-5 mortality data
18 available from UNICEF (<https://data.unicef.org/>). Future population projections (SSP2)
19 available from Wittgenstein Centre (<http://www.wittgensteincentre.org/dataexplorer>). Future
20 Global Burden of Disease mortality projection available from Institute for Health Metrics and
21 Evaluation ([http://ghdx.healthdata.org/record/ihme-data/global-life-expectancy-all-cause-
22 mortality-and-cause-specific-mortality-forecasts-2016-2040](http://ghdx.healthdata.org/record/ihme-data/global-life-expectancy-all-cause-mortality-and-cause-specific-mortality-forecasts-2016-2040)).

23 **Acknowledgements**

24 This work was supported by the Natural Environment Research Council (NERC) [grant
25 numbers NE/T013613/1, NE/T01363X/1]; Research Council of Norway (RCN) [grant
26 number 312601]; The Swedish Research Council for Health, Working Life and Welfare in
27 collaboration with the Swedish Research Council (Forte) [grant number 2019-01570]; and the
28 National Science Foundation (NSF) [grant number ICER-2028598]; coordinated through a

1 Belmont Forum partnership. Marsham was funded by HyCRISTAL, IMPALA and the NCAS
2 ACREW programme.

3 **References**

- 4 Arbuthnott, K., S. Hajat, C. Heaviside, and S. Vardoulakis. 2018. What is cold-related
5 mortality? A multi-disciplinary perspective to inform climate change impact
6 assessments. *Environment International* 121:119–129.
- 7 Azongo, D. K., T. Awine, G. Wak, F. N. Binka, and A. R. Oduro. 2012. A time series
8 analysis of weather variability and all-cause mortality in the Kasena-Nankana districts of
9 Northern Ghana, 1995-2010. *Global Health Action* 5:14–22.
- 10 Bastin, J. F., E. Clark, T. Elliott, S. Hart, J. Van Den Hoogen, I. Hordijk, H. Ma, S.
11 Majumder, G. Manoli, J. Maschler, L. Mo, D. Routh, K. Yu, C. M. Zohner, and T. W.
12 Crowther. 2019. Understanding climate change from a global analysis of city analogues.
13 *PLoS ONE* 14:1–13.
- 14 Basu, R., D. Pearson, L. Sie, and R. Broadwin. 2015. A Case-Crossover Study of
15 Temperature and Infant Mortality in California. *Paediatric and Perinatal Epidemiology*
16 29:407–415.
- 17 Berrang-Ford, L., A. J. Sietsma, M. Callaghan, J. C. Minx, P. F. D. Scheelbeek, N. R.
18 Haddaway, A. Haines, and A. D. Dangour. 2021. Systematic mapping of global research
19 on climate and health: a machine learning review. *The Lancet Planetary Health* 5:e514–
20 e525.
- 21 Chersich, M. F., M. D. Pham, A. Areal, M. M. Haghghi, A. Manyuchi, C. P. Swift, B.
22 Wernecke, M. Robinson, R. Hetem, M. Boeckmann, and S. Hajat. 2020. Associations
23 between high temperatures in pregnancy and risk of preterm birth, low birth weight, and
24 stillbirths: Systematic review and meta-analysis. *The BMJ* 371:1–13.
- 25 Chirombo, J., P. Ceccato, R. Lowe, D. J. Terlouw, M. C. Thomson, A. Gumbo, P. J. Diggle,
26 and J. M. Read. 2020. Childhood malaria case incidence in Malawi between 2004 and
27 2017: Spatio-temporal modelling of climate and non-climate factors. *Malaria Journal*
28 19:1–13.
- 29 Ebi, K. L., N. H. Ogden, J. C. Semenza, and A. Woodward. 2017. Detecting and Attributing
30 Health Burdens to Climate Change. *Environmental Health Perspectives*:1–8.

- 1 Egondi, T., C. Kyobutungi, S. Kovats, K. Muindi, R. Ettarh, and J. Rocklöv. 2012. Time-
2 series analysis of weather and mortality patterns in Nairobi's informal settlements.
3 *Global Health Action* 5:23–32.
- 4 Foreman, K. J., N. Marquez, A. Dolgert, K. Fukutaki, N. Fullman, M. McGaughey, M. A.
5 Pletcher, A. E. Smith, K. Tang, C. W. Yuan, J. C. Brown, J. Friedman, J. He, K. R.
6 Heuton, M. Holmberg, D. J. Patel, P. Reidy, A. Carter, K. Cercy, A. Chapin, D.
7 Douwes-Schultz, T. Frank, F. Goettsch, P. Y. Liu, V. Nandakumar, M. B. Reitsma, V.
8 Reuter, N. Sadat, R. J. D. Sorensen, V. Srinivasan, R. L. Updike, H. York, A. D. Lopez,
9 R. Lozano, S. S. Lim, A. H. Mokdad, S. E. Vollset, and C. J. L. Murray. 2018.
10 Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality
11 for 250 causes of death: reference and alternative scenarios for 2016–40 for 195
12 countries and territories. *The Lancet* 392:2052–2090.
- 13 Friedlingstein, P., R. M. Andrew, J. Rogelj, G. P. Peters, J. G. Canadell, R. Knutti, G.
14 Luderer, M. R. Raupach, M. Schaeffer, D. P. Van Vuuren, and C. Le Quéré. 2014.
15 Persistent growth of CO₂ emissions and implications for reaching climate targets.
16 *Nature Geoscience* 7:709–715.
- 17 Global Burden of Disease Collaborative Network. 2020. Global Burden of Disease Study
18 2019 (GBD 2019) Under-5 Mortality and Adult Mortality 1950-2019. Institute for
19 Health Metrics and Evaluation (IHME), Seattle, United States of America.
- 20 Green, H., J. Bailey, L. Schwarz, J. Vanos, K. Ebi, and T. Benmarhnia. 2019. Impact of heat
21 on mortality and morbidity in low and middle income countries: A review of the
22 epidemiological evidence and considerations for future research. *Environmental*
23 *Research* 171:80–91.
- 24 Das Gupta, P. 1993. *Decomposition of Rates: A User's Manual*. Standardization and
25 *Decomposition of Rates: A User's Manual*:P23-186.
- 26 Hajat, S., and T. Kosatky. 2010. Heat-related mortality: A review and exploration of
27 heterogeneity. *Journal of Epidemiology and Community Health* 64:753–760.
- 28 Hajat, S., S. Vardoulakis, C. Heaviside, and B. Eggen. 2014. Climate change effects on
29 human health: Projections of temperature-related mortality for the UK during the 2020s,
30 2050s and 2080s. *Journal of Epidemiology and Community Health* 68:641–648.

- 1 Harris, I. 2019. Release Notes for CRU TS v4.03: 15 May 2019.
2 https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.03/Release_Notes_CRU_TS4.03.txt.
- 3 Helldén, D., C. Andersson, M. Nilsson, K. L. Ebi, P. Friberg, and T. Alfvén. 2021. Climate
4 change and child health: a scoping review and an expanded conceptual framework. *The*
5 *Lancet Planetary Health* 5:e164–e175.
- 6 Honda, Y., M. Kondo, G. McGregor, H. Kim, Y. L. Guo, Y. Hijioka, M. Yoshikawa, K. Oka,
7 S. Takano, S. Hales, and R. S. Kovats. 2014. Heat-related mortality risk model for
8 climate change impact projection. *Environmental Health and Preventive Medicine*
9 19:56–63.
- 10 KC, S., and W. Lutz. 2017. The human core of the shared socioeconomic pathways:
11 Population scenarios by age, sex and level of education for all countries to 2100. *Global*
12 *Environmental Change* 42:181–192.
- 13 Kim, J., J. H. Kim, H. K. Cheong, H. Kim, Y. Honda, M. Ha, M. Hashizume, J. Kolam, and
14 K. Inape. 2016. Effect of climate factors on the childhood pneumonia in papua new
15 guinea: A time-series analysis. *International Journal of Environmental Research and*
16 *Public Health* 13:1–16.
- 17 Liu, T., C. Zhou, H. Zhang, B. Huang, Y. Xu, L. Lin, L. Wang, R. Hu, Z. Hou, Y. Xiao, J. Li,
18 X. Xu, D. Jin, M. Qin, Q. Zhao, W. Gong, P. Yin, Y. Xu, J. Hu, J. Xiao, W. Zeng, X. Li,
19 S. Chen, L. Guo, Z. Rong, Y. Zhang, C. Huang, Y. Du, Y. Guo, S. Rutherford, M. Yu,
20 M. Zhou, and W. Ma. 2021. Ambient Temperature and Years of Life Lost: A National
21 Study in China. *The Innovation* 2:100072.
- 22 Meinshausen, M., Z. R. J. Nicholls, J. Lewis, M. J. Gidden, E. Vogel, M. Freund, U. Beyerle,
23 C. Gessner, A. Nauels, N. Bauer, J. G. Canadell, J. S. Daniel, A. John, P. B. Krummel,
24 G. Luderer, N. Meinshausen, S. A. Montzka, P. J. Rayner, S. Reimann, S. J. Smith, M.
25 Van Den Berg, G. J. M. Velders, M. K. Vollmer, and R. H. J. Wang. 2020. The shared
26 socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to
27 2500. *Geoscientific Model Development* 13:3571–3605.
- 28 O’Neill, B. C., C. Tebaldi, D. P. Van Vuuren, V. Eyring, P. Friedlingstein, G. Hurtt, R.
29 Knutti, E. Kriegler, J. F. Lamarque, J. Lowe, G. A. Meehl, R. Moss, K. Riahi, and B. M.
30 Sanderson. 2016. The Scenario Model Intercomparison Project (ScenarioMIP) for
31 CMIP6. *Geoscientific Model Development* 9:3461–3482.

- 1 Phung, D., S. Rutherford, C. Chu, X. Wang, M. Nguyen, N. H. Nguyen, C. Manh, T. H.
2 Nguyen, and C. Huang. 2015. Temperature as a risk factor for hospitalisations among
3 young children in the Mekong Delta Area, Vietnam. *Occupational and Environmental*
4 *Medicine* 72:529–535.
- 5 Rocklöv, J., and K. L. Ebi. 2012. High Dose Extrapolation in Climate Change Projections of
6 Heat-Related Mortality. *Journal of Agricultural, Biological, and Environmental*
7 *Statistics* 17:461–475.
- 8 Romanello, M., A. McGushin, C. Di Napoli, P. Drummond, N. Hughes, L. Jamart, H.
9 Kennard, P. Lampard, B. Solano Rodriguez, N. Arnell, S. Ayeb-Karlsson, K. Belesova,
10 W. Cai, D. Campbell-Lendrum, S. Capstick, J. Chambers, L. Chu, L. Ciampi, C. Dalin,
11 N. Dasandi, S. Dasgupta, M. Davies, P. Dominguez-Salas, R. Dubrow, K. L. Ebi, M.
12 Eckelman, P. Ekins, L. E. Escobar, L. Georgeson, D. Grace, H. Graham, S. H. Gunther,
13 S. Hartinger, K. He, C. Heaviside, J. Hess, S. C. Hsu, S. Jankin, M. P. Jimenez, I.
14 Kelman, G. Kiesewetter, P. L. Kinney, T. Kjellstrom, D. Kniveton, J. K. W. Lee, B.
15 Lemke, Y. Liu, Z. Liu, M. Lott, R. Lowe, J. Martinez-Urtaza, M. Maslin, L. McAllister,
16 C. McMichael, Z. Mi, J. Milner, K. Minor, N. Mohajeri, M. Moradi-Lakeh, K.
17 Morrissey, S. Munzert, K. A. Murray, T. Neville, M. Nilsson, N. Obradovich, M. O.
18 Sewe, T. Oreszczyn, M. Otto, F. Owfi, O. Pearman, D. Pencheon, M. Rabhaniha, E.
19 Robinson, J. Rocklöv, R. N. Salas, J. C. Semenza, J. Sherman, L. Shi, M. Springmann,
20 M. Tabatabaei, J. Taylor, J. Trinanes, J. Shumake-Guillemot, B. Vu, F. Wagner, P.
21 Wilkinson, M. Winning, M. Yglesias, S. Zhang, P. Gong, H. Montgomery, A. Costello,
22 and I. Hamilton. 2021. The 2021 report of the Lancet Countdown on health and climate
23 change: code red for a healthy future. *The Lancet* 398:1619–1662.
- 24 Rylander, C., J. øyvind Odland, and T. M. Sandanger. 2013. Climate change and the potential
25 effects on maternal and pregnancy outcomes: An assessment of the most vulnerable - the
26 mother, fetus, and newborn child. *Global Health Action* 6.
- 27 Schinasi, L. H., J. R. Bloch, S. Melly, Y. Zhao, K. Moore, and A. J. de Roos. 2020. High
28 ambient temperature and infant mortality in Philadelphia, Pennsylvania: A case-
29 crossover study. *American Journal of Public Health* 110:189–195.
- 30 Schwalm, C. R., S. Glendon, and P. B. Duffy. 2020. RCP8.5 tracks cumulative CO2
31 emissions. *Proceedings of the National Academy of Sciences of the United States of*
32 *America* 117:19656–19657.

- 1 Sewe, M. O., A. Bunker, V. Ingole, T. Egondi, D. O. Astrom, D. M. Hondula, J. Rocklov,
2 and B. Schumann. 2018. Estimated Effect of Temperature on Years of Life Lost : A
3 Retrospective Time-Series Study of Low-, Middle-, and High-Income Regions.
4 *Environmental Health Perspectives* 126:1–12.
- 5 Sheffield, P. E., M. T. Herrera, E. J. Kinnee, and J. E. Clougherty. 2018. Not so little
6 differences: variation in hot weather risk to young children in New York City. *Public*
7 *Health* 161:119–126.
- 8 Sheffield, P. E., and P. J. Landrigan. 2011. Global climate change and children’s health:
9 Threats and strategies for prevention. *Environmental Health Perspectives* 119:291–298.
- 10 Smith, K. R., A. Woodward, D. Campbell-Lendrum, D. D. Chadee, Y. Honda, Q. Liu, J. M.
11 Olwoch, B. Revich, and R. Sauerborn. 2014. Human health impacts, adaptation and co-
12 benefits. Pages 709–754 in C. B. Field, V. R. Barros, D. J. Dokken, K. J. Mach, M. D.
13 Mastrandrea, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova, B.
14 Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea, and L. L. White,
15 editors. *Climate Change 2014 Impacts, Adaptation and Vulnerability: Part A: Global*
16 *and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report*
17 *of the Intergovernmental Panel on Climate Change.* Cambridge University Press,
18 Cambridge United Kingdom and New York, NY, USA.
- 19 Son, J. Y., J. T. Lee, and M. L. Bell. 2017. Is ambient temperature associated with risk of
20 infant mortality? A multi-city study in Korea. *Environmental Research* 158:748–752.
- 21 Spencer, S., T. Samateh, K. Wabnitz, S. Mayhew, H. Allen, and A. Bonell. 2022. The
22 Challenges of Working in the Heat Whilst Pregnant: Insights From Gambian Women
23 Farmers in the Face of Climate Change. *Frontiers in Public Health* 10:1–11.
- 24 Teutschbein, C., and J. Seibert. 2012. Bias correction of regional climate model simulations
25 for hydrological climate-change impact studies: Review and evaluation of different
26 methods. *Journal of Hydrology* 456–457:12–29.
- 27 Trisos, C. H., I. O. Adelekan, E. Totin, A. Ayanlade, J. Efitre, A. Gameda, K. Kalaba, C.
28 Lennard, C. Masao, Y. Mgaya, G. Ngaruiya, D. Olago, N. O. Simpson, and S.
29 Zakieldeen. 2022. Africa. Page 225 IPCC WGII Sixth Assessment Report.
- 30 UNICEF Data. 2019. UNICEF Data Warehouse. <https://data.unicef.org/>.

- 1 United Nations. 2015. The Millennium Development Goals Report. New York.
- 2 United Nations. 2019. World Population Prospects 2019, Online Edition, Rev.1. Department
3 of Economic and Social Affairs, Population Division.
- 4 United Nations. 2020. Goal 3: Ensure healthy lives and promote well-being for all at all ages.
5 <https://www.un.org/sustainabledevelopment/health/>.
- 6 Vicedo-Cabrera, A. M., N. Scovronick, F. Sera, D. Royé, R. Schneider, A. Tobias, C.
7 Astrom, Y. Guo, Y. Honda, D. M. Hondula, R. Abrutzky, S. Tong, M. de S. Z. S.
8 Coelho, P. H. N. Saldiva, E. Lavigne, P. M. Correa, N. V. Ortega, H. Kan, S. Osorio, J.
9 Kyselý, A. Urban, H. Orru, E. Indermitte, J. J. K. Jaakkola, N. Rytí, M. Pascal, A.
10 Schneider, K. Katsouyanni, E. Samoli, F. Mayvaneh, A. Entezari, P. Goodman, A. Zeka,
11 P. Michelozzi, F. De’Donato, M. Hashizume, B. Alahmad, M. H. Diaz, C. D. L. C.
12 Valencia, A. Overcenco, D. Houthuijs, C. Ameling, S. Rao, F. Di Ruscio, G. Carrasco-
13 Escobar, X. Seposo, S. Silva, J. Madureira, I. H. Holobaca, S. Fratianni, F. Acquotta,
14 H. Kim, W. Lee, C. Iniguez, B. Forsberg, M. S. Ragetti, Y. L. L. Guo, B. Y. Chen, S.
15 Li, B. Armstrong, A. Aleman, A. Zanobetti, J. Schwartz, T. N. Dang, D. V. Dung, N.
16 Gillett, A. Haines, M. Mengel, V. Huber, and A. Gasparrini. 2021. The burden of heat-
17 related mortality attributable to recent human-induced climate change. *Nature Climate*
18 *Change* 11.
- 19 Wang, Y., A. Wang, J. Zhai, H. Tao, T. Jiang, B. Su, J. Yang, G. Wang, Q. Liu, C. Gao, Z.
20 W. Kundzewicz, M. Zhan, Z. Feng, and T. Fischer. 2019. Tens of thousands additional
21 deaths annually in cities of China between 1.5 °C and 2.0 °C warming. *Nature*
22 *Communications* 10:1–7.
- 23 Wittgenstein Centre for Demography and Global Human Capital. 2018. Wittgenstein Centre
24 Data Explorer Version 2.0. <http://www.wittgensteincentre.org/dataexplorer>.
- 25 World Health Organization. 2005. Health and the Millennium Development Goals.
26 Switzerland.
- 27 World Health Organization. 2014. Quantitative risk assessment of the effects of climate
28 change on selected causes of death, 2030s and 2050s.
- 29 World Health Organization. 2018. Number of deaths in children aged <5, by cause.
30 <https://www.who.int/data/gho/data/indicators/indicator-details/GHO/number-of-deaths>.

1 WorldPop, and Center for International Earth Science Information Network (CIESIN). 2018.
2 WorldPop: Age and Sex structures. Columbia University.
3 <https://www.worldpop.org/geodata/listing?id=65>.

4 Xu, Z., R. A. Etzel, H. Su, C. Huang, Y. Guo, and S. Tong. 2012. Impact of ambient
5 temperature on children's health: A systematic review. *Environmental Research*
6 117:120–131.

7 Yé, Y., V. R. Louis, S. Simboro, and R. Sauerborn. 2007. Effect of meteorological factors on
8 clinical malaria risk among children: An assessment using village-based meteorological
9 stations and community-based parasitological survey. *BMC Public Health* 7:1–11.

10 Zhao, Q., Y. Guo, T. Ye, A. Gasparri, S. Tong, A. Overcenco, A. Urban, A. Schneider, A.
11 Entezari, A. M. Vicedo-Cabrera, A. Zanobetti, A. Analitis, A. Zeka, A. Tobias, B.
12 Nunes, B. Alahmad, B. Armstrong, B. Forsberg, S. C. Pan, C. Íñiguez, C. Ameling, C.
13 De la Cruz Valencia, C. Åström, D. Houthuijs, D. Van Dung, D. Royé, E. Indermitte, E.
14 Lavigne, F. Mayvaneh, F. Acquaotta, F. de' Donato, F. Di Ruscio, F. Sera, G. Carrasco-
15 Escobar, H. Kan, H. Orru, H. Kim, I. H. Holobaca, J. Kyselý, J. Madureira, J. Schwartz,
16 J. J. K. Jaakkola, K. Katsouyanni, M. Hurtado Diaz, M. S. Ragetti, M. Hashizume, M.
17 Pascal, M. de Sousa Zanotti Stagliorio Coêlho, N. Valdés Ortega, N. Rytí, N.
18 Scovronick, P. Michelozzi, P. Matus Correa, P. Goodman, P. H. Nascimento Saldiva, R.
19 Abrutzky, S. Osorio, S. Rao, S. Fratanni, T. N. Dang, V. Colistro, V. Huber, W. Lee, X.
20 Seposo, Y. Honda, Y. L. Guo, M. L. Bell, and S. Li. 2021. Global, regional, and national
21 burden of mortality associated with non-optimal ambient temperatures from 2000 to
22 2019: a three-stage modelling study. *The Lancet Planetary Health* 5:e415–e425.

23
24