

Summer convective precipitation over the UK and Europe from a regional weather projection.

Alan Gadian¹, Alan Blyth¹, Ralph Burton¹, Cindy Bruyere², James Done², James Groves¹, Greg Holland², Stephen Mobbs¹, Jutta Thielen-del-Pozo³, Mari Tye², James Warner⁴

¹NCAS, ICAS, University of Leeds, UK ²MMM, NCAR, USA, JRC, ³ISPRA, Italy ⁴University of Exeter, UK

Abstract:

Climate change caused by green house gas emissions is now following the trend of rapid warming consistent with a RCP8.5 forcing. Climate models are still unable to represent the mesoscale convective processes that occur at resolutions $\sim O(3\text{km})$ and are not capable of resolving precipitation patterns in time and space with sufficient accuracy to represent convection. In this poster, UK Met Office precipitation observations are compared with the simulations for the (1990-1995) period and a simulation of a near future period (2031-2036) for a regional nested weather model. (**Table 1**). The convection permitting model, resolution $\sim O(3\text{km})$, provides a good correspondence to the observational precipitation data and demonstrates the importance of explicit convection for future summer precipitation estimates. (**Table 2**). UK summer precipitation is reduced slightly ($\sim 10\%$) for (2031-2036) (**Figure 2**) and there is no evidence of an increase in the peak maximum hourly precipitation magnitude. A similar pattern is observed over the whole European inner model domain. The Kain-Fritsch convective parameterisation at a resolution $\sim O(12\text{km})$, over estimates the summer precipitation by $\sim 10\%$ for the UK compared with the $\sim O(3\text{km})$ simulations. In the future, the average precipitation rate per event increases (**Figure 3**), shorter heavier precipitation events increase (**Figure 4**), dry periods extend (**Figure 5**) and wet periods shorten. As part of the change, 10m winds of $< 3\text{ms}^{-1}$ become more common (**Figure 6**) – a scenario that would impact on power generation from wind turbines through calmer conditions causing more frequent pollution episodes.




Figure 2. Comparison of current and future precipitation patterns for the UK (JJA). Upper left panel indicates the UK precipitation amounts for summer (1990-1995) from the Centre of Ecology and Hydrology [CEH]. Upper right panel indicates the corresponding values from the high resolution $\sim O(3\text{km})$ driven by CESM data for (1990-1995). The lower panels show the model output data for years (2031-2036) driven by CESM data, for the inner and outer domains, with resolutions $\sim O(3\text{km})$ and $\sim O(12\text{km})$ respectively. All plots use the same colour scale.




Figure 3. Summer, (JJA), UK plots. Top row: changes in dry spell duration, for high resolution $\sim O(3\text{km})$, domain, D02, 2nd row: changes in heavy precipitation ($>7.6\text{mm}/\text{hour}$) for high resolution; the changes are produced by subtracting the (1990-1995) from the (2031-2036) average values at each pixel.




Figure 4. Summer, (JJA), UK plots. Intensity versus event duration plot for the difference in high resolution data $\sim O(3\text{km})$, domain, D02. Difference in precipitation intensity for (1990-1995) and (2031-2036); Changes are produced by subtracting the (1990-1995) from the (2031-2036) average values at each pixel.




Figure 5. Summer, (JJA): Change in dry spell event duration, for high resolution simulations. The changes are derived by subtracting the (1990-1995) from the (2031-2036) average values at each pixel.




Figure 1. Domain structure for the simulation. The outer domain (d01) resolution is 20km at $\pm 30^\circ$ and 8km at 68°N/S . The inner domain (d02) is one way nested at a ratio of 5:1

Figure 1 displays the domain structure used, with resolution from $< 3.2\text{km}$ and 2.0km at 35°N and 68°N in the inner domain, d02, increasing by a factor of five for the outer domain, implying resolutions $\sim O(3\text{km})$ and $\sim O(12\text{km})$. This covers 95% of the globe. 51 vertical stretched levels with a lid at 10mb. 1731 E-W grid points in both domains and 907 and 1001 grid points in the N - S directions in the outer and inner domains, respectively. CESM boundary conditions drive the simulations. Kain Fritsch, WSM6 microphysics, YSU boundary layer and IGBP-MODIS, a 4 layer Noah land surface schemes and CAM radiation schemes are used.




Figure 6. Summer, (JJA), changes in 10m wind and the change in the number of hours where the wind speed is less than 3ms^{-1} at 10m, using the high resolution $\sim O(3\text{km})$ simulations

Take home message: "For summer convective precipitation in future decades, there will be longer dry spells with shorter wet spells with more shorter but heavier wet events. Convectively permitting solutions capture ~ 10 times more than convectively parameterised simulations, with 20% increase in their total precipitation contribution; Europe will be 10% dryer"

Boundary Data	Domain / Source	Period of Simulation	Seasonal Average Precipitation
Era-Interim	d01 $\sim O(12\text{km})$	(1990-1995)	236 mm
Era-Interim	d02 $\sim O(3\text{km})$	(1990-1995)	221 mm
CESM version 1	d01 $\sim O(12\text{km})$	(1990-1995)	225 mm
CESM version 1	d02 $\sim O(3\text{km})$	(1990-1995)	204 mm
CEH observations	Met Off. data	(1990-1995)	205 mm
CESM version 1	d01 $\sim O(12\text{km})$	(2031-2036)	188 mm
CESM version 1	d02 $\sim O(3\text{km})$	(2031-2036)	182 mm

Table 1:

Six year average UK precipitation totals for summer months JJA for the control period (1990-1995) from the CEH observations and from the model simulations. Precipitation averages for the period (2031-2036) are also included for both domains.

Boundary Data	Domain / Source	Period of Simulation	Seasonal Average Precipitation
CESM version 1	d01 $\sim O(12\text{km})$	(1990-1995)	264 mm
CESM version 1	d02 $\sim O(3\text{km})$	(1990-1995)	196 mm
CESM version 1	d01 $\sim O(12\text{km})$	(2031-2036)	254 mm
CESM version 1	d02 $\sim O(3\text{km})$	(2031-2036)	177 mm

Table 2:

Six year average European domain precipitation average totals for summer months JJA for the control period (1990-1995) and for the period (2031-2036), for the low and high resolution simulations.

Summary and Key Points.

This is an analysis of convective precipitation over the whole European domain, (the inner model domain) which large enough to enable several day cycles, using time steps that can explicitly resolve air motions. The results from the high resolution convection permitting simulation $\sim O(3\text{km})$ suggests that the lower resolution K-F convective parameterisation $\sim O(12\text{km})$ simulations do not always replicate the convective changes and produce a larger effective precipitation efficiency. The UK only results for (1990-1995) are in good agreement with the six year observed precipitation totals with the convection permitting simulations. For the future simulation (2031-2036), especially in the south east of the UK, the results suggest that the summers will possibly be drier, with longer dry spells, shorter wet spells and heavier precipitation. Overall, the UK will be $\sim 10\%$ dryer for the summer periods assessed. The K-F convection parameterisation simulations underestimates the importance of short, (< 4 hours) heavy precipitation ($>7.6\text{mm}/\text{hour}$) events, which is more apparent in the future time period. The convection permitting solutions capture about ~ 10 times as many of these short heavy precipitation events, and both convection approaches suggest a $\sim 20\%$ increase in their contribution to precipitation by (2031-2036). The average precipitation intensity per event increases. The UK domain results are mirrored in the European domain. With longer drier periods and shorter wet events suggested, the results are possibly more pronounced. The increase in average precipitation for each event is even more apparent over the European domain. This will result in more flash floods over Europe in restricted catchments areas and have implications for agriculture. Associated with this is an average increase of approximately 150 hours in the occurrence of 10m summer wind speeds of less than 3ms^{-1} .

Acknowledgements.

The authors acknowledge the support of NERC funding for NCAS (National Centre for Atmospheric Sciences), the "leadership award" from EPSRC / NERC for the use of the ARCHER CRAY XC30 supercomputer, the support of the UK's national high-performance computing service, and for NCAR (National Center for Atmospheric Research) support and allocation of supercomputer time under the ECEP project and for JRC, ISPRA assistance.